



Small clusters of fermions

J. Carbonell, R. Lazauskas

► To cite this version:

J. Carbonell, R. Lazauskas. Small clusters of fermions. 17th International IUPAP Conference on Few-Body Problems in Physics Few-Body 17, Jun 2003, Durham, United States. pp.S106-S108. in2p3-00022254v2

HAL Id: in2p3-00022254

<https://hal.in2p3.fr/in2p3-00022254v2>

Submitted on 2 Dec 2004

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


Small clusters of fermions

J. Carbonell and R. Jazauskas^a

^aLab. Physique Subatomique et Cosmologie, 53 Av. Martyrs, 38026 Grenoble, France

Several theoretical studies [1] indicate that there is no any reasonable chance for $3n$ and $4n$ to be bound. GANIL experiment [2] suggesting tetra-neutron has not been confirmed but these no-result had however the merit of pushing theoreticians at work to clarify the possible existence of small n -clusters. If the situation seems clear for $A=3$ and 4 , it is less well established for a larger number of neutrons. The search of multi-neutron resonances raises also some interest and a recent experiment scanning the $4n$ continuum in the $d(^8\text{He}, ^6\text{Li})4n$ reaction reports a 2-3 MeV width structure [3].

In this issue it is enlightening to make a parallel with a similar, better-known, fermionic system: the ^3He atomic clusters. Since small ^3He droplets can exist [4], should we expect some stability islands (see figure below) in the continuum neutron states going from $N=2$ to $N=\infty$. If yes, where? If not, why?



The two-body interactions of these systems look at first glance quite similar. We have compared in Fig. 1 the n - n AV18 [5] and the ^3He - ^3He Aziz [6] S-wave potentials. They have been rescaled by the corresponding masses $\frac{M}{\hbar^2}$ and the resulting length units, as well as the inter-particle distance d are fm and \AA respectively. Corresponding low energy parameters are given in Table 1. In the n - n case we have also considered Reid93 [7] and MT13 [8] potentials, the latter being adjusted to reproduce the experimental scattering length. None of these system supports a bound dimer ($a < 0$) but ^3He seems less favorable

Table 1

Low energy n - n and ^3He - ^3He parameters.

	n-n(fm)				He-He(\AA)
	Av18	Reid	MT13	Exp.	Aziz 91
a	-18.49	-17.54	-18.59	-18.59 ± 0.4	-7.24
r_0		2.85	2.94	2.75 ± 0.1	13.5
η_c	1.08	1.09	1.10		1.30

to make clusters. This can be seen by calculating the critical values of the scaling factor η_c introduced in the potential $V^{(\eta)}(r) = \eta V_{nn}(r)$, which bounds a dimer. For n - n , this value is around $\eta_c = 1.08$ whereas for He-He is sensibly greater $\eta_c = 1.30$ (see Table 1).




Figure 1. Comparison between n - n and ${}^3\text{He}$ - ${}^3\text{He}$ potentials.




Figure 2. Critical value of the scaling factor for n (circles) and ${}^3\text{He}$ (squares)

When examining larger systems we first consider the bosonic case. Despite the absence of dimers, "bosonic" neutron trimers and tetramers do exist with binding energies $B_{n_3} \approx 1$ MeV and $B_{n_4} \approx 10$ MeV. This is however not the case for atomic ${}^3\text{He}$, suggesting once more that neutron clusters should be favored with respect to atomic ${}^3\text{He}$. They all disappear when the Pauli principle is imposed, but can reappear – as in ${}^3\text{He}$ – if the number of interacting particles is increased. The existence of small clusters results thus from a compromise between an attractive pairwise interaction and the effective Pauli repulsion.

In order to study such a balance, we have investigated in several directions [9]: *(i)* scaling factor in V_{nn} , *(ii)* three-neutron interactions (TnI), *(iii)* influence of n - n P- waves *(iv)* confining the system in an harmonic oscillator (HO) trap and *(iv)* "dimer"-"dimer" scattering. It is always possible to bind 3- and 4- n states by modifying the usual n - n and/or TnI models but the violation has to be very strong, producing serious anomalies. Thus, in case *(i)* one needs a large scaling factor $\eta_c \sim 3$. Using ad-hoc TnI, one gets a very compact object in which NN force becomes even repulsive. Keeping the usual S-wave n - n potential, the enhancement of n - n P-waves required to bound a 3- or 4- n system is such that they become themselves resonant!

We have confined $N=2,3,4$ n in an HO trap with fixed frequency ω and size parameter $b = \sqrt{\frac{\hbar}{m\omega}}$. HO is the only external field in which "internal" and "center of mass" energies can be properly separated. In absence of n - n forces the "internal" energies are known analytically but can be obtained as well by solving the N -body "internal" problem, with pairwise HO potential of frequency $\left(\frac{\omega}{\sqrt{N}}\right)$. The effect of n - n interaction has been evaluated by solving the internal problem with

$$V_{ij} = \frac{1}{2} m \left(\frac{\omega}{\sqrt{N}} \right)^2 r_{ij}^2 + V_{nn}(r_{ij})$$

and calculating the difference between the pure HO energy and the $HO + V_{nn}$ one: $B_N = E_{HO}^{(N)} - E_{HO+nn}^{(N)}$.

Results concerning the ground state are given in Table 2 for several values of b . Some comments are in order: *(i)* there is a clear indication of paring effect when going from

Table 2

Binding energies B_N of N neutrons in an HO trap with size parameter b .

N	J^π	$b=2$				$b=3$				$b=4$			
		$E_{HO}^{(N)}$	B_N	$\frac{B_N}{N}$	$\frac{B_N}{E_{HO}}$	$E_{HO}^{(N)}$	B_N	$\frac{B_N}{N}$	$\frac{B_N}{E_{HO}}$	$E_{HO}^{(N)}$	B_N	$\frac{B_N}{N}$	$\frac{B_N}{E_{HO}}$
2	0^+	15.55	6.34	3.17	0.41	6.91	3.13	1.56	0.45	3.89	1.81	0.93	0.47
3	$\frac{3}{2}^-$	41.47	9.74	3.25	0.23	18.43	4.41	1.47	0.24	10.36	2.55	0.85	0.25
4	0^+	67.39	15.30	3.58	0.23	29.95	7.40	1.69	0.25	16.82	4.31	1.08	0.26

$N=2 \rightarrow 3 \rightarrow 4$ (ii) one has always $B_4 > 2B_2$, suggesting an effective attraction between dineutrons (iii) the binding energy per particle increases when going from $N=2$ to $N=4$ (iv) the ratio $\frac{B_N}{E_{HO}}$ tends to a constant value independent of b . The preceding results tend to indicate that there is a benefit when going from 2 to 4. The main difference in respect to ${}^3\text{He}$ has been found in the role of P-waves. Their influence in n case is attractive but very small whereas they significantly contribute to the ${}^3\text{He}$ binding energy ($\sim 40\%$). The reason for such a different behaviour is the hard core radius of the corresponding potentials, which differ by a factor 3 (see Fig. 1). The centrifugal barrier is one order of magnitude smaller in ${}^3\text{He}$ and the effective potential is, contrary to n case, still attractive in regions where it can play a role. This difference can be dramatic in binding larger fermion systems.

Despite the negative results quoted in [1], the question of larger neutron clusters merits some attention. At present, the strongest argument against their existence are the mean-field results, all concluding to an unbounded infinite nuclear matter but it should be possible to reach the same conclusion "from below", i.e. from a systematic study of few-neutron systems. This can be performed by studying the N -dependence of the critical scaling factor η_c , calculated simultaneously for n and ${}^3\text{He}$. Our technology allow us to reach only $N=4$ with the results displayed on Fig. 2. The value $\eta_c^{(n)}$ makes a large jump when passing from $N=2$ to $N=3$ but starts to decreases when going from $N=3$ to $N=4$. Is that a pure numerical accident or, as in ${}^3\text{He}$, an indication of a descent towards the $\eta = 1$ axis?. More powerful methods could go far beyond $N=4$ and draw a definite conclusion on this problem.

REFERENCES

1. S.C. Pieper, Phys. Rev. Lett. **90** (2003) 252501; contrib. to FB17 and refs. therein
2. F.M. Marques et al, Phys. Rev. **C65**, 044006 (2002).
3. D. Baumel, private communication
4. R. Guardiola, J. Navarro, Phys. Rev. Lett. **84**, 1144 (2001).
5. R.B. Wiringa, V.G.J. Stoks, R. Schiavilla, Phys. Rev. **C51** (1995) 38
6. R.A. Aziz and M.J. Slaman, J. Chem. Phys. **94** (1991) 8047
7. V.G. Stoks, R.A. Klomp, C.P. Terheggen, J.J. de Swart, Phys. Rev. **C49** (1994) 2950
8. R.A. Malfliet, J. Tjon, Nucl. Phys. **A127** (1969) 161
9. R. Lazauska, PhD. Université de Grenoble (2003)